



AMN gene

amnion associated transmembrane protein

Normal Function

The *AMN* gene provides instructions for making a protein called amnionless. This protein is involved in the uptake of vitamin B12 (also called cobalamin) from food. Vitamin B12, which cannot be made in the body and can only be obtained from food, is essential for the formation of DNA and proteins, the production of cellular energy, and the breakdown of fats. This vitamin is involved in the formation of red blood cells and maintenance of the brain and spinal cord (central nervous system).

The amnionless protein is primarily found embedded in the outer membrane of kidney cells and cells that line the small intestine. Amnionless attaches (binds) to another protein called cubilin, anchoring cubilin to the cell membrane. Cubilin can interact with molecules and proteins passing through the intestine or kidneys. During digestion, vitamin B12 is released from food. As the vitamin passes through the small intestine, cubilin binds to it. Amnionless helps transfer the cubilin-vitamin B12 complex into the intestinal cell. From there, the vitamin is released into the blood and transported throughout the body. In the kidneys, amnionless and cubilin are involved in the reabsorption of certain proteins that would otherwise be released in urine.

Health Conditions Related to Genetic Changes

Imerslund-Gräsbeck syndrome


At least 30 mutations in the *AMN* gene have been found to cause Imerslund-Gräsbeck syndrome. This condition is characterized by low levels of vitamin B12 in the body, which leads to a blood disorder known as megaloblastic anemia. About half of affected individuals also have excess protein in their urine (proteinuria), and some have neurological problems.

AMN gene mutations that cause Imerslund-Gräsbeck syndrome reduce the amount or function of the amnionless protein. Without amnionless acting as an anchor, cubilin is not attached to cells in the small intestine or kidneys and cannot bind to vitamin B12 and other molecules and proteins needed in the body. As a result, instead of being taken into intestinal cells, vitamin B12 is released from the body. A shortage of this essential vitamin impairs the proper development of red blood cells, leading to megaloblastic anemia. Low levels of vitamin B12 can also affect the central nervous system, causing neurological problems. In addition, without amnionless function in the kidneys, proteins are not reabsorbed into the body and are instead released in the urine, leading to proteinuria.

Chromosomal Location

Cytogenetic Location: 14q32.32, which is the long (q) arm of chromosome 14 at position 32.32

Molecular Location: base pairs 102,922,430 to 102,930,842 on chromosome 14 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- amnionless
- amnionless homolog
- PRO1028
- protein amnionless
- protein amnionless precursor
- visceral endoderm-specific type 1 transmembrane protein

Additional Information & Resources

Educational Resources

- Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline: Vitamin B12
<https://www.ncbi.nlm.nih.gov/books/NBK114302/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28AMN%5BTIAB%5D%29+OR+%28amnionless%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- AMNIONLESS, MOUSE, HOMOLOG OF
<http://omim.org/entry/605799>

Research Resources

- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=AMN%5Bgene%5D>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=14604
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/81693>
- UniProt
<http://www.uniprot.org/uniprot/Q9BXJ7>

Sources for This Summary

- OMIM: AMNIONLESS, MOUSE, HOMOLOG OF
<http://omim.org/entry/605799>
- Beech CM, Liyanarachchi S, Shah NP, Sturm AC, Sadiq MF, de la Chapelle A, Tanner SM. Ancient founder mutation is responsible for Imerslund-Gräsbeck Syndrome among diverse ethnicities. *Orphanet J Rare Dis*. 2011 Nov 13;6:74. doi: 10.1186/1750-1172-6-74.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22078000>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3226546/>
- Fyfe JC, Madsen M, Højrup P, Christensen EI, Tanner SM, de la Chapelle A, He Q, Moestrup SK. The functional cobalamin (vitamin B12)-intrinsic factor receptor is a novel complex of cubilin and amnionless. *Blood*. 2004 Mar 1;103(5):1573-9. Epub 2003 Oct 23.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14576052>
- Kozyraki R, Cases O. Vitamin B12 absorption: mammalian physiology and acquired and inherited disorders. *Biochimie*. 2013 May;95(5):1002-7. doi: 10.1016/j.biochi.2012.11.004. Epub 2012 Nov 20. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23178706>
- Namour F, Dobrowoljski G, Chery C, Audonnet S, Feillet F, Sperl W, Gueant JL. Luminal expression of cubilin is impaired in Imerslund-Grasbeck syndrome with compound AMN mutations in intron 3 and exon 7. *Haematologica*. 2011 Nov;96(11):1715-9. doi: 10.3324/haematol.2011.043984. Epub 2011 Jul 12.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21750092>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3208692/>
- Watkins D, Rosenblatt DS. Lessons in biology from patients with inborn errors of vitamin B12 metabolism. *Biochimie*. 2013 May;95(5):1019-22. doi: 10.1016/j.biochi.2013.01.013. Epub 2013 Feb 10. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23402785>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/AMN>

Reviewed: April 2014

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services